Saskatchewan Agriculture and Food

Crop Report

Published by Saskatchewan Agriculture and Food
ISSN 0701 7065

Report Number 18, August 7, 2007

Crop Report for August 5, 2007

Farmers have four per cent of the 2007 crop combined, according to Saskatchewan Agriculture and Food's weekly crop report.

A further nine per cent has either been swathed or is ready to be straight combined. There has been some harvest activity in all regions of the province, and in all crops.

Fall cereals are reported to be about 80 per cent in good to excellent condition. Spring cereals are reported to be between 41 to 61 per cent in good to excellent condition. Oilseed crops are reported to be 57 to 60 per cent in good to excellent condition. Pulse crops are 49 to 58 per cent in good to excellent condition.

Topsoil moisture conditions on crop, hay, and pasture land deteriorated in all regions during the past week. Over three-quarters of the province's crop land is reported to be in short or very short topsoil moisture condition.

Desiccation of pulse crops is underway. While many areas of the province are looking for a rain shower to help finish off crops, reporters in some areas indicated that rain will no longer be beneficial and may reduce crop quality.

First-cut haying operations are nearing completion, with 89 per cent baled or put into silage. Although some areas have begun a second cut, with 12 per cent completed provincially, many reporters do not expect a second cut of hay in their areas.

Hot, dry conditions were the main source of crop damage during the past week. Many crop reporters indicated that yields will be reduced due to the heat. Insect damage came from bertha armyworms, wheat stem maggot, wheat midge, aphids, grasshoppers, flea beetles, diamond back moth larvae, and sawflies. Disease pressures came from rust, root rot, aster yellows, sclerotinia, glume blotch, ascochyta blight, alternaria black spot, and mildew. Other sources of crop damage during the past week included hail, wind, and gophers.

1 Year Ago

Four per cent of the 2006 crop was combined, and seven per cent had been swathed or was ready to be straight-combined.

First-cut haying operations were nearing completion with 95 per cent baled or put into silage.

Hot, dry conditions were the main sources of crop damage.

Saskatchewan Harvest Progress at August 5, 2007

	% In Swath or Ready to Str.Cmb.		Average	% Combined		Average
	2007	2006	2002-06	2007	2006	2002-06
Winter wheat	22	28	16	21	47	16
Spring wheat	3	2	1	0	1	0
Durum	7	5	1	0	3	1
Oats	2	3	1	1	0	0
Barley	4	4	2	1	1	0
Fall rye	28	23	26	36	57	21
Triticale	11	15	6	1	12	3
Flax	0	0	0	0	0	0
Canola	12	8	3	0	0	0
Mustard	21	13	5	4	4	1
Sunflowers	3	0	0	0	0	0
Lentils	29	23	8	16	17	4
Peas	21	20	9	26	20	7
Canary seed	1	1	0	0	0	0
Chickpeas	5	6	2	2	4	1
Total	**9**	**7**	**3**	**4**	**4**	**1**

For further information contact:
Terry Bedard, P.Ag.
Agricultural Economist
Policy Branch
Saskatchewan Agriculture and Food
306-787-5956
e-mail: policybranch@agr.gov.sk.ca

Available on the SAF Internet site at: www.agr.gov.sk.ca

Please let us know if you no longer need to receive a printed copy.

Agriculture and Agri-Food Canada | Agriculture et Agroalimentaire Canada

CANADA – SASKATCHEWAN Crop Insurance

South eastern Saskatchewan (Crop Districts 1, 2, & 3ase)

The south eastern region was generally hot and dry during the past week with a few scattered showers. An average of 1 mm of precipitation was reported, with an average high of 2 mm reported in Crop District 1b. Heat and drought stress were reported across the region. Wind damage was reported in CDs 1a, 2b, and 3ase. In the Odessa area, the wind scattered and twisted pea swaths. The wind lodged crops in the Zehner area.

Five per cent of the 2007 south eastern crop has been combined and a further 11% has been swathed or is ready to be straight-combined. Forty-four per cent of the winter wheat, 41% of the fall rye, 31% of the peas, and 16% of the lentils have been combined. Some barley has been combined in the Wawota area and yielded about 60 bu/acre – farmers had expected 80 bu/acre earlier in the season. In the Ceylon area, the peas were running between 30 to 35 bu/acre.

Crop conditions continue to deteriorate with the hot, dry weather pushing a higher percentage of crops into the fair and some to the poor categories.

In the Lampman area, the winter wheat only went about 75% of what was expected. Also in that area, farmers are worried about light weight oats and barley. In the Kisbey area, there has been some bleaching in wheat swaths.

Topsoil moisture conditions on cropland deteriorated this past week. Thirty per cent of the crop land is rated as having very short topsoil moisture conditions, compared with 17% last week. Conditions also deteriorated on hay and pasture land with 33% of the land rated as very short, compared to 19% last week.

Insect damage came from aphids, bertha armyworms, wheat midge, wheat stem maggots, diamondback moth larvae, flea beetles, saw flies, and grasshoppers. Some insecticide spraying was done. Stag head was reported in canola in the Lampman area, and root rot was reported in the Grenfell area. Wheat stem maggot is reducing yields by 10% or more in the Lampman and Kennedy areas.

First-cut haying operations are almost complete with 91% baled or silaged, and 7% lying in the swath. Quality is generally expected to be good. Second-cut haying has begun in some areas with 28% baled or put into silage and 10% cut across the region. Many reporters do not expect a second cut in their area.

Spring crops are being desiccated. Cattle in the Minton area are suffering from the heat and the flies.

South western Saskatchewan (CDs 3asw, 3an, 3b, & 4)

The past week was again hot and dry in the southwest, with some strong winds. An average of 3 mm of precipitation was reported during the past week, with an average high of 9 mm reported in CD 3bs. Heat and drought stress were reported across the region. Wind damage was reported in CDs 3an, 3bs, 3bn, and 4b. Swaths were blown around in the Moose Jaw and Eyebrow areas. Some canola, lentil, and pea crops were shattered in the Glenbain area. Crops were lodged in the Elrose area. Hail damage was reported in the Admiral, Elrose, and Eastend areas.

Eleven per cent of the 2007 south western crop has been combined and a further 18% is swathed or is ready to be straight-combined. Thirty-eight per cent of the winter wheat, 51% of the fall rye, 51% of the peas, and 28% of the lentils have been combined.

Wheat samples are shriveled in the Fife Lake area. In the Limerick area, crops are dropping leaves and turning pale in colour. In the Mossbank area, peas were harvested at 86 to 89 days. In the Hazenmore area, early reported yields include peas at 18 bu/acre, lentils at 10 bu/acre, and mustard at 7 bu/acre. Early harvested samples in the Vanguard area indicate very light bushel weights. Red lentil yields in the Tyner area varied from 750 to 1500 lbs/acre. Yellow peas are maturing green in the Gull Lake area and grading as feed. In the Sceptre area, yellow peas came in at 30 bu/acre.

Crop conditions in the southwest continue to deteriorate as the majority of the spring crops fall

into the fair and poorer categories.

Topsoil moisture conditions on crop land remain similar to last week, with 59% of the land rated in very short topsoil moisture condition. For hay and pasture land, 70% is rated as having very short topsoil moisture.

Insect pressures came from grasshoppers, flea beetles, and sawflies. Disease pressures came from rust. Gophers continue to damage crops in many areas. It is hoped that the recent approval of liquid strychnine will help in reducing populations in the worst regions.

Haying operations are winding down in the southwest, with 96% of the first-cut hay baled or put into silage. Quality is expected to be good. Most reporters do not expect to see a second cut in their area.

Rain in some areas on the Monday holiday helped reduce the fire hazard somewhat. Farmers will still need to be prepared for fires while harvesting.

Pastures are getting very dry. Water sources are low for livestock in the Stewart Valley and Shaunavon areas.

East central Saskatchewan (Crop Districts 5 & 6a)

The east central region was hot with some thunderstorms during the past week. An average of 4 mm of precipitation was reported, with an average of 8 mm reported in CD 5a. Drought, heat, and wind stress were reported across the region. Crops were lodged in the Insinger area from the wind and the rain. In the Bethune area, the wind scattered canola and lentil swaths. Hail damage was reported in several areas. In the Saltcoats area, hail damage was mostly on the canola. In the Ituna area, the hail wiped out crops from 5 to 100%, and caused damage to power lines and buildings. Large hail was reported in the Liberty area with a lot of damage.

One per cent of the 2007 east central crop has been combined and 5% has been swathed or is ready to be straight-combined. Eight per cent of the winter wheat, 17% of the fall rye, 4% of the lentils, and 4% of the peas have been combined. Desiccation is underway.

Early yield reports from the Holdfast area show winter wheat at 30 to 50 bu/acre and peas at 30 to 40 bu/acre. Winter wheat in the Semans area went about 40 bu/acre.

Over 50% of crops remain categorized as being in good to excellent condition, but the heat is moving more of them into the fair and poor categories.

Topsoil moisture conditions deteriorated, with 36% of the crop land rated as having adequate topsoil moisture, compared with 54% last week. Conditions also deteriorated on hay and pasture land with 42% of the land rated as having adequate topsoil moisture, compared to 52% last week.

Insect damage came from bertha armyworms, sawflies, wheat midge, grasshoppers, aphids, wheat stem maggot, and diamondback moth larvae. Some insecticide application is underway. Disease pressure came from leaf disease, rust, aster yellows, stag head, and smut.

First-cut haying operations are progressing, with 90% baled or silaged, and a further 7% lying in the swath. Quality is expected to be good. The second cut is just getting underway with 5% baled or made into silage. Many reporters do not expect to see a second cut in their area. Hay yields were reported to be down 20 to 30% in the Goodeve area.

West central Saskatchewan (Crop Districts 6b & 7)

The west central region saw warm, dry weather this past week, with some isolated showers. An average of 6 mm of precipitation was reported during the past week, with an average of 10 mm reported in CD 7a. Heat and drought stress caused damage across the region. Crops are turning white in several areas. Wind damage was reported in several areas in CDs 6b and 7b. Hail damage was reported in the Imperial and Hanley areas. Hail in the southern third of RM 283 caused damage of up to 100%.

One per cent of the 2007 west central crop has been combined and a further 7% has been swathed or is ready to be straight-combined. Sixteen per cent of the fall rye, 15%

of the winter wheat, 6% of the lentils and 3% of the peas have been combined. Pulse desiccation is underway.

In the Rosetown area, fall rye was reported to yield around 40 bu/acre.

Over 50% of the crops remain in good to excellent condition, but the percentage in the fair and poor categories is increasing.

Topsoil moisture conditions deteriorated, with 23% of the crop, hay and pasture land rated in very short moisture condition, compared with 14% last week. Pastures are in poor condition.

Insect pressures came from bertha armyworms, flea beetles, sawflies, diamondback moth larvae, and wheat midge. Disease pressures came from sclerotinia, glume blotch, alternaria black spot, and ascochyta blight.

Cutting and baling the 2007 hay crop is winding down with 96% baled or silaged, and 3% lying in the swath. Quality is generally expected to be good. Nineteen per cent of the second cut crop has been baled and a further 44% cut, though a number of reporters do not expect to harvest a second cut.

North eastern Saskatchewan (Crop Districts 8 & 9ae)

The north eastern region was sunny, warm, and dry during the past week. An average of 3 mm of precipitation was reported during the past week, with an average of 10 mm reported in CD 9ae. Heat and drought stress were reported in CD 8. Rainfall and cooler temperatures are needed for crops. Wind damaged crops in the Arborfield area.

One per cent of 2007 crop has been swathed or is ready to straight combine in the northeast. No combining has been done yet. Peas are starting to be desiccated.

Over 50% of the crops remain in good to excellent condition, though crops are deteriorating and moving into the fair and poor categories.

In the Porcupine Plain area, it is estimated that later-seeded cereal crops may have lost 50% of their yield because of the heat. Crops are starting to burn in the Tisdale area and yield losses are expected, especially in canola.

Topsoil moisture conditions continued to deteriorate this past week, with 42% of the crop, hay and pasture land rated as having short or very short topsoil moisture conditions, compared with 15 to 20% last week.

Insect damage came from bertha armyworms, wheat midge, leaf mites, and diamondback moth larvae. Insecticide spraying continues.

Haying operations are coming along with close to 90% of the 2007 crop baled or silaged. Quality is expected to be good. Harvesting a second cut has begun with 5% baled or silaged and a further 14% lying in the swath. Many reporters do not expect a second cut in their area.

North western Saskatchewan (Crop Districts 9aw & 9b)

The north western region had another hot, dry week with some hail. An average of 3 mm of precipitation was reported, with an average of 4 mm reported in CD 9aw. Heat and drought stress were reported across the region. A few good showers would help crops. Wind damaged crops in the Speers and North Battleford areas. Hail was reported in the Meota area. Golf-ball sized hail was reported in the North Battleford area on Aug. 6.

Four per cent of the 2007 north western crop has been swathed or is ready to be straight combined. One per cent of each of the oats and the barley, and 13% of the fall rye have been made into greenfeed.

For the most part, the majority of the crop remains in the good to excellent categories, though larger percentages are being rated as fair and/or poor.

Early-seeded crops are burning in the Hafford area. Crops are also burning in the Meota area. Peas are wilting off in the Speers area. Last week's heat did more crop damage in the Leask area than the heat of the previous weeks. Harvest is coming in fast in the Lloydminster area and crop prospects are poor.

Topsoil moisture conditions deteriorated during the past week, with 16% of the crop land and 22% of the hay and pasture land rated as having adequate topsoil moisture, compared to 38% and 36% respectively last week. Pastures are starting to turn brown in the Barthel area.

Bertha armyworms, diamondback moth larvae, and grasshoppers all caused crop damage during the past week. Insecticide application is underway. Sclerotinia was reported in the Shellbrook area. Gophers caused damage in the St. Walburg area.

Eighty-six per cent of the hay crop has been baled or silaged, and a further 9% is lying in the swath. Quality is expected to be good. Many reporters do not expect to see a second cut in their area.

Saskatchewan Crop Conditions, Aug 5, 2007

	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oat	Barley	Triticale	
% excellent	18	20	8	6	11	8	1	
% good	61	60	46	35	50	48	53	
% fair	18	19	36	44	32	36	43	
% poor	2	1	9	13	6	7	3	
% very poor	1	0	1	2	1	1	0	
	Flax	Canola	Mustard	Sun-flower	Lentil	Pea	Canary Seed	Chick-pea
% excellent	8	8	7	9	9	9	5	13
% good	52	49	37	80	48	49	54	36
% fair	35	34	31	10	35	32	34	40
% poor	5	8	21	1	7	9	6	10
% very poor	0	1	4	0	1	1	1	1

Southeast Saskatchewan Crop Conditions, Aug 5, 2007

	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oat	Barley	Triticale	
% excellent	21	29	9	6	10	8	0	
% good	64	56	49	43	51	48	90	
% fair	14	13	37	44	32	38	10	
% poor	1	2	5	7	7	6	0	
% very poor	0	0	0	0	0	0	0	
	Flax	Canola	Mustard	Sun-flower	Lentil	Pea	Canary Seed	Chick-pea
% excellent	7	5	4	10	5	14	2	3
% good	52	53	52	87	56	59	66	58
% fair	36	34	29	3	38	26	30	37
% poor	5	7	15	0	1	1	2	2
% very poor	0	1	0	0	0	0	0	0

Southwest Saskatchewan Crop Conditions, Aug 5, 2007

	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oat	Barley	Triticale	
% excellent	19	18	7	6	6	6	0	
% good	44	66	18	21	16	25	43	
% fair	32	16	49	49	42	47	54	
% poor	3	0	22	20	30	16	3	
% very poor	2	0	4	4	6	6	0	
	Flax	Canola	Mustard	Sun-flower	Lentil	Pea	Canary Seed	Chick-pea
% excellent	9	8	8	20	6	5	11	17
% good	25	23	21	0	26	35	32	25
% fair	37	48	28	49	47	42	39	43
% poor	28	18	33	31	17	16	12	13
% very poor	1	3	10	0	4	2	6	2

East Central Saskatchewan Crop Conditions, Aug 5, 2007

	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oat	Barley	Triticale	
% excellent	19	21	11	10	13	12	0	
% good	68	44	56	46	55	51	35	
% fair	12	33	28	32	28	32	65	
% poor	1	2	5	12	4	5	0	
% very poor	0	0	0	0	0	0	0	
	Flax	Canola	Mustard	Sun-flower	Lentil	Pea	Canary Seed	Chick-pea
% excellent	10	10	6	0	10	11	8	0
% good	57	50	50	80	56	57	43	43
% fair	31	33	36	20	31	29	47	57
% poor	2	5	8	0	3	3	2	0
% very poor	0	2	0	0	0	0	0	0

West Central Saskatchewan Crop Conditions, Aug 5, 2007

	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oat	Barley	Triticale	
% excellent	9	11	7	6	10	8	9	
% good	88	72	58	64	58	58	50	
% fair	3	16	31	28	31	30	41	
% poor	0	1	4	2	1	3	0	
% very poor	0	0	0	0	0	1	0	
	Flax	Canola	Mustard	Sun-flower	Lentil	Pea	Canary Seed	Chick-pea
% excellent	10	9	8	0	14	14	2	6
% good	54	52	47	0	61	60	59	77
% fair	36	30	36	100	22	23	30	17
% poor	0	8	9	0	2	2	9	0
% very poor	0	1	0	0	1	1	0	0

Northeast Saskatchewan Crop Conditions, Aug 5, 2007

	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oat	Barley	Triticale	
% excellent	17	22	9	0	15	6	0	
% good	50	42	50	58	47	60	80	
% fair	28	36	39	40	37	32	20	
% poor	4	0	1	2	1	2	0	
% very poor	1	0	1	0	0	0	0	
	Flax	Canola	Mustard	Sun-flower	Lentil	Pea	Canary Seed	Chick-pea
% excellent	0	7	0	n/a	0	7	10	0
% good	54	54	67		65	64	52	75
% fair	45	33	33		27	21	38	25
% poor	1	5	0		8	7	0	0
% very poor	0	1	0		0	1	0	0

Northwest Saskatchewan Crop Conditions, Aug 5, 2007

	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oat	Barley	Triticale	
% excellent	9	0	5	0	8	5	0	
% good	85	27	43	100	53	35	94	
% fair	6	48	37	0	27	46	0	
% poor	0	25	15	0	12	14	6	
% very poor	0	0	0	0	0	0	0	
	Flax	Canola	Mustard	Sun-flower	Lentil	Pea	Canary Seed	Chick-pea
% excellent	13	5	0	n/a	0	9	0	0
% good	36	41	28		100	47	100	100
% fair	50	38	0		0	31	0	0
% poor	1	16	72		0	13	0	0
% very poor	0	0	0		0	0	0	0

Weekly Rainfall Summary

(in millimeters)

for the week ending August 4, 2007

Crop Dist.	No.	Municipality Name	Past Week	Since April 1
1A	1	Argyle	NIL	194
	2	Mount Pleasant	NIL	265*
	3	Enniskillen	NIL	265
	33 A	Moose Creek	NIL	220
	33 B	Moose Creek	1	221
	34	Browning	NIL	277
	35	Benson	NIL	197
	63 A	Moose Mountain	NIL	306
	64	Brock	3	273
	65	Tecumseh	NIL	219
1B	91	Maryfield	NIL	256
	92 B	Walpole	2	337
	93	Wawken	NIL	289
	94	Hazelwood	NIL	336
	95	Golden West	NIL	278
	123	Silverwood	3	297
	124	Kingsley	2	379
	125	Chester	2	240
	151 B	Rocanville	13	331
	153	Willowdale	1	301
	155	Wolseley	TR	274
2A	36	Cymri	NIL	307
	66	Griffin	NIL	214
	67	Weyburn	NIL	246
	68	Brokenshell	NIL	269
	69	Norton	NIL	227
	96	Fillmore	4	250
	97	Wellington	TR	174
	99	Caledonia	3	269
2B	126	Montmartre	5	197
	127	Francis	TR	185
	128	Lajord	NIL	136
	129	Bratt's Lake	2	154
	131	Baildon	3	196
	156 B	Indian Head	1	234
	157	South Qu'appelle	NIL	273
	158	Edenwold	0	179
	160	Pense	NIL	174
	161	Moose Jaw	2	185
	191	Marquis	NIL	169
3ASE	9	Surprise Valley	NIL	269
	38 A	Laurier	NIL	261
	38 B	Laurier	NIL	249
	39 A	The Gap	NIL	218
	39 B	The Gap	NIL	245
3ASW	10	Happy Valley	NIL	303
	12	Poplar Valley	NIL	205
	40	Bengough	NIL	266
	43 A	Old Post	NIL	203
	43 B	Old Post	NIL	172
	44	Waverley	NIL	177
	70	Key West	NIL	201
	71 A	Excel	NIL	252
	73 A	Stonehenge	1	150
	73 B	Stonehenge	NIL	180
3AN	101 B	Terrell	NIL	182
	102	Lake Johnston	NIL	166
	103	Sutton	NIL	204*
	132	Hillsborough	6	142
	193	Eyebrow	NIL	184
	194	Enfield	NIL	147
3BS	45	Mankota	10	156
	75 A	Pinto Creek	13	165
	75 B	Pinto Creek	3	153*
	76 A	Auvergne	6	136
	76 B	Auvergne	18	184
	77	Wise Creek	14	225
	78 A	Grassy Creek	6	227
	78 B	Grassy Creek	17	198
	78 C	Grassy Creek	6	238
	105	Glenbain	TR	210
	106	Whiska Creek	7	176
	108	Bone Creek	5	174
3BN	136	Coulee	NIL	155
	138	Webb	NIL	171
	165	Morse	NIL	73
	166	Excelsior	NIL	105
	167	Sask. Landing	NIL	171
	225	Canaan	NIL	112
	226	Victory	NIL	168
	228 A	Lacadena	NIL	173
	228 B	Lacadena	NIL	178
	257	Monet	14	162*
4A	49 A	White Valley	NIL	155
	49 B	White Valley	NIL	224
	51	Reno	NIL	170
	79	Arlington	NIL	214
	109 A	Carmichael	5	174
	110	Piapot	NIL	155
	111 A	Maple Creek	NIL	138
	111 B	Maple Creek	NIL	155
4B	139	Gull Lake	NIL	162
	141	Big Stick	NIL	198
	142	Enterprise	NIL	235
	230	Clinworth	NIL	138
	231	Happyland	NIL	210
	232	Deer Forks	NIL	154
5A	183	Fertile Belt	5	267
	185	Mcleod	NIL	255
	186	Abernethy	NIL	199
	213	Saltcoats	8	242
	216 A	Tullymet	4	249
	216 B	Tullymet	NIL	207
	241	Calder	8	193
	243	Wallace	25	200*
	245 A	Garry	20	278
	245 B	Garry	24	229
	246	Ituna Bon Accord	10	231
	247	Kellross	35	238
	248	Touchwood	10	134
5B	271	Cote	1	222
	273	Sliding Hills	7	210
	275 A	Insinger	14	270
	276	Foam Lake	13	290
	277	Emerald	1	195
	307	Elfros	NIL	264
	308	Big Quill	TR	210
	335	Hazel Dell	5	268
	336	Sasman	NIL	231
	337	Lakeview	NIL	241
	338	Lakeside	NIL	233
	366 B	Kelvington	TR	195
	367 B	Ponass Lake	NIL	193
6A	189	Lumsden	TR	198
	190 A	Dufferin	2	129
	190 B	Dufferin	2	144
	190 C	Dufferin	NIL	172
	219 B	Longlaketon	1	133
	219 C	Longlaketon	9	152
	220	Mckillop	NIL	133
	221	Sarnia	NIL	171
	222	Craik	NIL	172
	251	Big Arm	30	188
	252	Arm River	NIL	135
	279 A	Mount Hope	NIL	118
	279 B	Mount Hope	8	173
	280	Wreford	NIL	134
	309	Prairie Rose	NIL	232
	312	Morris	NIL	132
	313	Lost River	NIL	143
	339 B	Leroy	0	210
	340 A	Wolverine	NIL	218
	340 B	Wolverine	3	211
	343	Blucher	1	170
6B	254	Loreburn	1	145
	283	Rosedale	4	179
	284	Rudy	20	194
	285	Fertile Valley	7	196
	286	Milden	12	183
	314	Dundurn	NIL	191
	344	Corman Park	NIL	249
	345	Vanscoy	NIL	239
	346	Perdue	17	243
	376 A	Eagle Creek	2	223
	376 B	Eagle Creek	15	207
	403	Rosthern	3	229
	404	Laird	NIL	187
7A	287	St. Andrews	11	160
	288	Pleasant Valley	13	182
	292	Milton	NIL	205
	317 A	Marriott	8	230
	317 B	Marriott	14	219
	318 A	Mountain View	18	178
	318 B	Mountain View	5	212
	320	Oakdale	27	220
	321	Prairiedale	15	231
7B	347 A	Biggar	NIL	216
	350 A	Mariposa	TR	230
	350 B	Mariposa	1	187
	351	Progress	10	252
	352	Heart's Hill	NIL	213
	377	Glenside	10	231
	378 A	Rosemount	5	219
	378 B	Rosemount	5	207
	379	Reford	TR	196
	380	Tramping Lake	3	189
	382	Eye Hill	10	315
	409	Buffalo	7	271
	410	Round Valley	NIL	210
8A	395	Porcupine	NIL	261
	397	Barrier Valley	1	238
	398	Pleasantdale	5	284
	428	Star City	2	250
	456	Arborfield	3	281
	457	Connaught	1	274
	458	Willow Creek	5	240
	486	Moose Range	2	179
	487	Nipawin	7	267
8B	369	St. Peter	NIL	257
	370	Humboldt	1	219
	371	Bayne	2	180
	372	Grant	2	166
	429 B	Flett's Springs	3	261
	431	St. Louis	7	290
9AE	488	Torch River	8	256
	520	Paddockwood	10	216
9AW	406	Mayfield	5	216
	435	Redberry	10	269
	436	Douglas	6	216
	463	Duck Lake	3	274
	464	Leask	3	253
	466	Meeting Lake	4	198
	467	Round Hill	TR	176
	493	Shellbrook	5	221
	494	Canwood	1	220
	496	Spiritwood	3	228
9B	438	Battle River	TR	193*
	440	Hillsdale	1	175
	442	Manitou Lake	NIL	210
	468	Meota	TR	165
	498 A	Parkdale	NIL	183
	498 B	Parkdale	NIL	164
	499	Mervin	1	217
	501 A	Frenchman Butte	TR	186
	501 B	Frenchman Butte	NIL	187
	502	Britannia	3	218
	561	Loon Lake	4	216
	588 B	Meadow Lake	5	199
	588 C	Meadow Lake	TR	207*
	622	Beaver River	3	220

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R.M.

* incomplete

Municipality No: A, B, C, and D - more than one reporter

Cumulative Rainfall
From: April 1, 2007
To: August 4, 2007



Cumulative Rainfall
From: April 1, 2007
To: August 4, 2007



Crop Land Topsoil Moisture conditions
August 5, 2007



Crop Land Topsoil Moisture conditions
August 5, 2007



Crop Districts
and
Rural Municipalities